



Influence of N, P, K, Rhizobium and Vermicompost on Soil Health Parameters and Yield of Green Gram (*Vigna radiata* L.) var. Samrat

Tanubrata Kapat ^{a++*}, Arun Alfred David ^{a#}, Joy Dawson ^{bt},
Tarence Thomas ^{at}, Neha Toppo ^{a‡}, Bidisha Chowdhury ^{a++},
Paritosh Sen ^{a++} and Raghvendra Kumar Kushwaha ^{a++}

^a Department of Soil Science and Agricultural Chemistry, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj – 211 007 (Uttar Pradesh), India.

^b Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj – 211 007 (Uttar Pradesh), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i74731>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/117786>

Original Research Article

Received: 28/03/2024

Accepted: 03/06/2024

Published: 06/06/2024

ABSTRACT

An experiment was conducted on "Influence of N, P, K, Rhizobium and Vermicompost on Soil Health Parameters and Yield of Green Gram (*Vigna radiata* L.) var. Samrat". during *zaid* season 2023-2024 at the central research farm, department research farm of Soil Science and Agricultural Chemistry, The design applied was 3x3 RBD having three levels each of vermicompost and N.P.K

⁺⁺ M.Sc. Student;

[#] Associate Professor;

[†] Professor;

[‡] Ph. D Scholar;

*Corresponding author: E-mail: kapattanubrata@gmail.com;

Cite as: Kapat, Tanubrata, Arun Alfred David, Joy Dawson, Tarence Thomas, Neha Toppo, Bidisha Chowdhury, Paritosh Sen, and Raghvendra Kumar Kushwaha. 2024. "Influence of N, P, K, Rhizobium and Vermicompost on Soil Health Parameters and Yield of Green Gram (*Vigna Radiata* L.) Var. Samrat". *International Journal of Plant & Soil Science* 36 (7):282-89. <https://doi.org/10.9734/ijpss/2024/v36i74731>.

i.e., 0, 50, and 100% respectively and three levels of rhizobium treated seed @ 0, 50, 100%. The result obtained in treatment T9 (N25P50K25 kg ha⁻¹ + vermicompost @ 5 t ha⁻¹ and rhizobium @ 200g 10 kg⁻¹) has showed a slight change in soil pH *i.e.*, 6.79 and EC *i.e.*, 0.208 dS m⁻¹. The post-harvest soil samples have showed significant increase in pore space, water holding capacity and organic carbon *i.e.*, about 49.20 %, 47.59 % and 0.49 % respectively. Whereas available N, P and K found to be 334.12 kg ha⁻¹, 34.36 kg ha⁻¹, and 204.28 kg ha⁻¹ respectively. The increase in NPK (kg ha⁻¹) was found to be significant among all treatments in green gram cultivation and soil quality improvement. The best results among all treatment combination with respect to plant height (65.76 cm), number of leaves plant⁻¹ (44.34), number of branches plant⁻¹ (14.58), number of grain pod⁻¹ (14.28), number of pod plant⁻¹ (42.11) and seed yield q ha⁻¹ (16.67) were found in treatment combination T9 (N25P50K25 kg ha⁻¹ + vermicompost @ 5 t ha⁻¹ and Rhizobium @ 200g 10 kg⁻¹). T9 gave the highest gross return with ₹ 1,49,030 and the benefit of ₹ 29,849.13 with highest cost benefit ratio *i.e.*, 1:1.38. vermicompost along with other combinations has showed improvement in soil health as well as in crop development and yield. Study has also revealed that the application of N, P, and K with vermicompost was an excellent source for fertilization than sole fertilizers.

Keywords: Soil health; green gram; vermicompost; rhizobium; NPK; yield etc.

1. INTRODUCTION

The term "soil health" describes the capability of soil to function. Profitability of farms, sustainable crop production, water and nutrient cycling, and climate regulation all depend on healthy soil. Evaluation of soil quality in relation to land production and management sustainability within the framework of soil functions are the two main goals of soil health assessment. A number of characteristics of soil have been found to be markers of soil health. Researchers have been assessing the health of the soil in research plots using a variety of soil health management techniques Manther M. Tahat [1].

Green gram is popularly known as "Moong Dal." "It is widely cultivated throughout Asia, including India. Pulses are the main source of protein particularly for vegetarians and contribute about 14 % of the total protein of average Indian diet". "The per capita availability of pulses in India has been continuously decreasing which is 32.52 g day⁻¹ against the minimum requirement of 80 g day⁻¹ per capita prescribed by Indian Council of Medical Research (ICMR). Therefore, agricultural scientists must to evolve strategy to increase production of pulses to meet the protein requirements of increasing population of the country" (K. Sowmya, 2016).

"Vermicompost is relatively resistant to microbial degradation. However, it is essential for establishing and maintaining the optimum soil physical condition for plant growth. It is a good source of N for sustainable crop production, but its availability remains an important issue due to its bulky nature, while inorganic fertilizer is no longer affordable to poor farmers due to its high cost.

Vermicompost contains nutrients especially nitrogen, phosphorous, potassium that are easily taken up by plants for fast growth. These composts provide a source of all necessary macro and micro-nutrients in available forms, thereby improving the physical and biological properties of the soil" [2].

"Nitrogen can be fixed by Green Gram from the atmosphere; it reacts to a tiny initial dose of nitrogenous fertilizers. The best results have been obtained when applying 15–25 kg N ha⁻¹. Increasing the nitrogen dose could potentially hinder nitrogen fixation capacity by diminishing nodule formation and growth. Nitrogen is indispensable for all crops, playing a crucial role in enhancing protein content and overall nutritional yield. Inadequate nutrient levels can result in stunted growth and a yellowish-green hue in plants. Moreover, nitrogen accelerates the growth and development of living tissues, particularly boosting tiller count in cereals and enhancing the photosynthetic activity of green plants" [3].

"Phosphorus plays a crucial role in boosting pulse productivity. It exerts a notably beneficial influence on the formation of nodules, root development, and nitrogen fixation in legume crops. The fixation process in legume crops, reliant on atmospheric nitrogen, demands considerable energy, with phosphorus being essential to fulfill its ATP (adenosine triphosphate) requirements" [4].

"It is better for the environment to use biofertilizers. Biofertilizers can be highly useful in fixing atmospheric nitrogen since they enhance and boost soil fertility. In India, the

most important type of bio-fertilizers are rhizobium inoculants that are specifically made for leguminous crops found that most of the agricultural contribution of biological nitrogen fixation comes from the symbiotic connection between legumes and rhizobium species” [5].

2. METHODOLOGY

The detailed treatment combination is showed in Table 1. and field experiment had been conducted during the Zaid season 2022 central research farm of department of Soil Science and Agricultural Chemistry, located at 25°24'30" North latitude 81°51'10" East longitude and 98m above mean sea level. Representing the Agro-ecological sub region [North Alluvium plain zone (0- 1% slope)] and Agro-climatic zone (Upper Gangetic plain region). “Argo climatically, The maximum temperature of the location ranges between 46°C and seldom falls below 4°C-5°C. The relative humidity ranges between 20-94%. The average rainfall of this area is around 1100mm annually”. The soil samples were randomly collected from one site in the experiment plot prior to tillage operation from a depth of 0-15 cm. The volume of the soil sample then reduced by conning and quartering the composites soil samples. Following this, they underwent air drying and were sieved through a 2 mm mesh, preparing them for both physical and chemical analysis. Soil physical analysis was conducted after post-harvest operations. After 60 days crop harvest soil sample was collected from field. Physical properties textural class, soil colour, bulk density $Mg\ m^{-3}$, particle density $Mg\ m^{-3}$, pore space %, water holding capacity % were analyzed. Soil chemical analysis is done after post-harvest operations were following, pH, EC $dS\ m^{-1}$, organic carbon %, available N $kg\ ha^{-1}$, P $kg\ ha^{-1}$, K $kg\ ha^{-1}$. The trial was laid out in a randomized block design with three replications; plot size was 2 x 2 m for crop seed rate is 20 $kg\ ha^{-1}$ moong.

3. RESULTS AND DISCUSSION

3.1 Soil Parameters

The composition of N, P, K and vermicompost have significant increase on the soil parameters. The increase of pore space %, water holding capacity %, organic carbon, available nitrogen, phosphorus, potassium, with the improvement of soil parameters, Table 2. Revealed that application of different level s of Vermicompost and N, P, K have following roll, on soil. In treatment T1 lowest data observed (Due to no application of N,P,K, Vermicompost and Rhizobium) pore space 42.31 and 40.30%, water holding capacity 41.15 and 40.45% and T9 shows the highest (Due to highest application of N,P,K, Vermicompost and Rhizobium) pore space 49.20% and 49.21%, water holding capacity 47.59% and 47.65% at 0-15cm and 15- 30cm depth of soil respectively. Also, in Table 3. Shown highest in bulk density T1 is $1.31Mg\ m^{-3}$ and 1.32 and particle density $2.39Mg\ m^{-3}$ and $2.40Mg\ m^{-3}$ (Because no application of Vermicompost in this treatment), with lowest in T9 bulk density $1.18Mg\ m^{-3}$ and $1.20Mg\ m^{-3}$ and partical density is $2.25Mg\ m^{-3}$ and $2.26Mg\ m^{-3}$ (Because highest application of Vermicompost in this treatment) respectively in 0- 15cm and 15-30 cm depth of soil. Table 4. shown that in Treatment T1 have highest pH 7.24 and 7.34 , EC 0.28 and 0.29 $dS\ m^{-1}$, organic carbon 0.37 and 0.33%, nitrogen 296.5 and 288.5 $kg\ ha^{-1}$, phosphorus 23.32 and 23.33 $kg\ ha^{-1}$, Potassium 134.81 and 132.41 $kg\ h^{-1}$ (Due to no application of N,P,K, Vermicompost and Rhizobium)and T9 have lowest pH 6.79 and 6.82, EC 0.19 and 0.20 $dS\ m^{-1}$, organic carbon 0.50% and 0.49% , nitrogen 334.12 and 326.12 $kg\ ha^{-1}$, Phosphorus 34.36 and 34.89 $kg\ ha^{-1}$, potassium 204.28 and 200.61 $kg\ ha^{-1}$ respectively(Due to highest application of N,P,K, Vermicompost and Rhizobium) in 0-15cm and 15-30cm depth of soil.

Table 1. Methodology employed for analysis

Particulars	Method employed
Bulk density ($Mg\ m^{-3}$)	Graduated Measuring Cylinder [6]
Partical density ($Mg\ m^{-3}$)	Graduated Measuring Cylinder [6]
Pore space(%)	Graduated Measuring Cylinder [6]
Soil pH (1:2)	pH meter (Glass electrode)
Soil EC ($dS\ m^{-1}$)	EC meter (Conductivity Bridge)
Organic Carbon (%)	Wet Oxidation Method [7]
Available Nitrogem ($Kg\ ha^{-1}$)	Kjeldhal Method [8]
Available Phosphorus ($Kg\ ha^{-1}$)	Colorimetric method [9]
Available Potassium ($Kg\ ha^{-1}$)	Flame photometric method [10]

Table 2. Treatment combination of green gram

S. No.	Treatment combination	Symbol
T ₁	(@N ₀ P ₀ K ₀ kg ha ⁻¹ +Vermicompost @ 0 t ha ⁻¹ and Rhizobium @ 0g 10 kg ⁻¹ Seed)	I1V1R1
T ₂	(@N ₀ P ₀ K ₀ kg ha ⁻¹ +Vermicompost @ 2.5t ha ⁻¹ and Rhizobium @ 100g 10 kg ⁻¹ Seed)	I1V2R2
T ₃	(@N ₀ P ₀ K ₀ kg ha ⁻¹ +Vermicompost @ 5 t ha ⁻¹ and Rhizobium @ 200g 10 kg ⁻¹ Seed)	I1V3R3
T ₄	(@N _{12.5} P ₂₅ K _{12.5} kg ha ⁻¹ +Vermicompost@ 0 t ha ⁻¹ and Rhizobium @ 0g 10 kg ⁻¹ Seed)	I2V1R1
T ₅	(@N _{12.5} P ₂₅ K _{12.5} Kg ha ⁻¹ +Vermicompost @ 2.5 t ha ⁻¹ and Rhizobium @ 100g 10 kg ⁻¹ Seed),	I2V2R2
T ₆	(@N _{12.5} P ₂₅ K _{12.5} kg ha ⁻¹ +Vermicompost @ 5 t ha ⁻¹ and Rhizobium @ 200g 10 kg ⁻¹ Seed),	I2V3R3
T ₇	(@N ₂₅ P ₅₀ K ₂₅ kg ha ⁻¹ +Vermicompost@ 0 t ha ⁻¹ and Rhizobium @ 0g 10 kg ⁻¹ Seed)	I3V1R1
T ₈	(@N ₂₅ P ₅₀ K ₂₅ kg ha ⁻¹ +Vermicompost @ 2.5 t ha ⁻¹ and Rhizobium @ 100g 10 kg ⁻¹ Seed),	I3V2R2
T ₉	(@N ₂₅ P ₅₀ K ₂₅ kg ha ⁻¹ +Vermicompost @ 5 t ha ⁻¹ and Rhizobium @ 200g 10 kg ⁻¹ Seed).	I3V3R3

Table 3. Effect of different levels of N, P, K, Vermicompost and Rhizobium on physical properties of soil

Treatments	Bulk density (Mg m ⁻³)		Particle density (Mg m ⁻³)		Pore space (%)		Water holding capacity (%)	
	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm
T ₁	1.31	1.32	2.39	2.40	42.31	42.30	41.15	40.45
T ₂	1.29	1.30	2.39	2.39	43.70	43.69	40.91	41.22
T ₃	1.19	1.21	2.32	2.33	44.12	44.18	42.08	42.74
T ₄	1.26	1.27	2.34	2.35	45.89	45.87	43.48	44.63
T ₅	1.23	1.26	2.31	2.31	46.65	46.66	44.01	44.78
T ₆	1.23	1.25	2.24	2.25	46.12	46.13	44.42	45.63
T ₇	1.22	1.25	2.30	2.30	46.31	46.31	45.06	46.27
T ₈	1.30	1.31	2.28	2.28	48.65	48.66	46.28	47.15
T ₉	1.18	1.20	2.25	2.26	49.20	49.21	47.59	47.65
F-test	S	S	S	S	S	S	S	S
S. Em. (±)	0.05	0.05	0.03	0.03	0.03	0.03	0.8	1.25
C.D.@5 %	0.02	0.02	0.01	0.01	0.14	0.14	0.38	0.59

Table 4. Effect of different levels of N, P, K, Vermicompost and Rhizobium on chemical properties of soil

Treatments	pH		EC (dS m ⁻¹)		Organic Carbon (%)		Nitrogen (kg ha ⁻¹)		Phosphorus (kg ha ⁻¹)		Potassium (kg ha ⁻¹)	
	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm
T1	7.24	7.34	0.28	0.29	0.37	0.33	296.50	288.50	23.32	23.33	134.81	132.41
T2	7.40	7.50	0.27	0.28	0.40	0.35	301.45	295.45	25.36	25.17	149.38	148.01
T3	7.42	7.52	0.25	0.26	0.42	0.36	306.14	296.14	26.04	25.58	154.73	154.39
T4	7.28	7.38	0.22	0.23	0.45	0.39	314.67	307.00	27.24	27.31	155.94	155.87
T5	7.05	7.11	0.23	0.24	0.44	0.39	314.51	309.84	28.42	28.56	165.24	162.24
T6	7.02	7.08	0.23	0.24	0.48	0.43	318.45	316.49	30.30	29.71	173.45	168.78
T7	6.91	7.01	0.22	0.23	0.46	0.43	320.20	319.50	31.15	31.40	186.65	179.65
T8	6.89	6.92	0.21	0.22	0.48	0.46	325.07	316.07	33.75	33.01	196.98	186.98
T9	6.79	6.82	0.19	0.20	0.50	0.49	334.12	326.12	34.36	33.89	204.28	200.61
F-test	S	S	S	S	S	S	S	S	S	S	S	S
S. Em.(±)	0.33	0.36	0.02	0.01	0.03	0.05	3.07	3.07	0.91	1	4.35	4.38
C.D.@ 5%	0.16	0.17	0.01	0.02	0.06	0.02	1.45	1.45	0.43	0.47	2.05	2.07

The T₉ treatment is found to be best, followed by T₈ and T₇, as depicted by their respective physical and chemical properties. The application of N, P, K, vermicompost, and rhizobium eventually proves to have a positive impact on the soil, helping to maintain its fertility. T₁ demonstrates the least effect on soil parameters among the treatments [11-14]

3.2 Economy of Green Gram Crop

According to following Table 5. The economy of different treatment concerned, the treatment T₉ provides highest net profit of ₹29849.13 with cost benefit ratio 1:1.34 however, the minimum net profit of ₹10095 was recorded in the treatment T₁ with cost benefit ratio is 1:1.8 [15-18].




Fig. 1. Effect of different levels of pH, EC(dS m⁻¹) and Organic Carbon(%) in soil




Fig. 2. Effect of different levels of Nitrogen (kg ha⁻¹), Phosphorous(kg ha⁻¹) and Potassium (kg ha⁻¹) in soil

Table 5. Effect of different treatment combination on cost benefit ratio (C: B) of green gram

Treatment	Yield (q ha ⁻¹)	Selling price (₹ q ⁻¹)	Gross return (₹ ha ⁻¹)	Total cost of cultivation (₹ ha ⁻¹)	Net Profit (₹ ha ⁻¹)	Cost Benefit ratio (C:B)
T1	7.33	9000	65970	55875.00	10095	1:1.18
T2	7.67	9000	69030	56275.00	12755	1:1.22
T3	9.33	9000	83970	66275.00	17695	1:1.26
T4	11.33	9000	101970	88727.97	13242.03	1:1.14
T5	11.67	9000	105030	88727.97	16302.03	1:1.18
T6	12.67	9000	114030	99954.40	14075.6	1:1.14
T7	13.33	9000	119970	99954.40	20015.6	1:1.20
T8	14.00	9000	126000	101180.87	24819.13	1:1.24
T9	16.67	9000	149030	111180.87	29849.13	1:1.34

4. CONCLUSION

It revealed from trial that the various level of N,P,K, vermicompost and rhizobium seed treatment used in the experiment, the treatment combination T9 [N25P50K25 kg ha⁻¹+Vermicompost @ 5 t ha⁻¹ and Rhizobium @ 200g 10 kg⁻¹ Seed] was found to be the best treatment with highest gross return of ₹ 149030 which gave benefit of ₹29849.13 with highest cost benefit ratio 1:3.4 for green gram. Therefore, T9 can be recommended for profitable production of green gram (*Vigna radiata* L.) var. Samrat and it is also found that treatment T9 has shown significance results over soil physical and chemical properties. Therefore, Combination of N, P, K, vermicompost and rhizobium treated seed is found to be better among all treatments for both soil health and green gram production.

ACKNOWLEDGEMENT

During my academic journey and research, my advisor and the Head of the Department of Soil Science and Agricultural Chemistry at NAI, SHUATS, have been valuable sources of support.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Manther M. Tahat. Soil Health and Sustainable Agriculture Albert MH. Munsell soil color charts. Munsell Color Company Inc., Baltimore; 1954. Bouyoucos GJ. The Hydrometer as a new method for the mechanical analysis of soils. Soil Science. 2020;23:343-353.
- Manisha Singh. Effect of vermicompost and panchagavya on growth and yield of greengram (*Vigna radiata* L.); 2022.
- Patel BK. Effect of various sources of nitrogen and phosphorus on growth, yield and economics of summer green Gram (*Vigna radiata* L. Wilczek); 2020.
- Sipai AH, Jat JR, Rathore BS. Effect of phosphorus, sulphur and biofertilizer on growth, yield and nodulation in mungbean on loamy sand soils of Kutch. A Scitechnol Journal. 2016;51(1):51-56.
- Sneha M. Isolation and Characterization of Rhizobium from Green Gram (*Vigna radiata*); 2022.
- Muthuvel P, Udayasoorian C, Natesan R, Ramaswami PR. Introduction to soil analysis. Tamil Nadu Agricultural University, Coimbatore; 1992.
- Walkley A, Black IA. An examination of degtjareff method for determining soil organic matter, and proposed modification of the chromic acid titration method. Soil Science. 1934;37:29- Wilcox RR. STDs: Definition. A Textbook of STDs, 1950;11.
- Subbiah BV, Asija GL. A rapid procedure for the estimation of available nitrogen in soils. Current Science. 1956;25:259-260.
- Olsen SR, Cole CV, et al. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. USDA Circular 939. US Government Printing Office, Washington DC; 1954.
- Toth SJ, Prince AL. Estimation of cation exchange capacity and exchangeable calcium, potassium, and sodium contents of soils by flame photometer techniques. Soil Sci. 1949;67:439-445.

11. Abhishek Singh Yadav, Tarence Thomas, Arun Alfred David. Effect of different organic manures on yield and physico-chemical properties of soil by kharif maize (*Zea mays* L.) on eastern plain zone of Uttar Pradesh of Prayagraj region. Int. J. Curr. Microbiol. App. Sci. 2020;9(9):634-640.
12. K. Sowmya. 2023 Productivity and profitability of green gram [*Vigna radiata* (L.) Wilczek] under system of crop intensification.
13. Jackson ML. Soil chemical analysis. Prentice-Hall Inc., Englewood Cliffs, NJ, 1958:498. Manvender Singh Khangaroat, Tarence Thomas, Arun Alfred David. Effect of different level of NPK and biochar on soil physico - chemical properties, yield and attributes of green gram (*Vigna radiata* L.) Var. – RMG 975 (KeshwanandMung 1). International Journal of Chemical Studies. 2020;8(5): 1302-1307.
14. Muhammad M, Kutawa AB, Adamu M. Influence of NPK fertilizer and poultry manure on the growth of okra (*Abelmoschus esculentus* L. Moench) in Northern Sudan Savanna region of Nigeria. International Journal of Horticulture, Agriculture and Food science (IJHAF). 2020;4(6).
15. Pathak K, Kalita MK, Barman U, Hazarika BN, Saha NN. Response of summer green gram (*Vigna radiata*) to inoculation and nitrogen levels in Barak valley zone of Assam. Annals of Agri. Res. 2001;22:123-124.
16. Inuwa NM, Fagge AA, Mohamed AML, Lado A, Aliyu AM. Effect of Stand Densities and NPK Fertilizer on Growth and Calyx Yield of Roselle (*Hibiscus sabdariffa* L.) in Sudan Savanna Zone of Nigeria: An Approach towards Growth Characters. J. Exp. Agric. Int. 2022 Sep. 1;44(10):238-47. Available:<https://journaljeai.com/index.php/JEAI/article/view/2020>
17. Jannat SA, Haque MA, Nahar K, Ali MKJ, Chowhan S, Hossian MM, Rahman MH. Integrated effect of vermicompost and chemical fertilizers on the yield & yield contributing characteristics of brinjal (*Solanum melongena* L.). Asian J. Adv. Agric. Res. 2023 Dec. 8;23(4):39-43. Available:<https://journalajaar.com/index.php/AJAAR/article/view/477>
18. Joshi R, Vig AP, Singh J. Vermicompost as soil supplement to enhance growth, yield and quality of *Triticum aestivum* L.: A field study. International Journal of Recycling of Organic Waste in Agriculture. 2013 Dec;2:1-7.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/117786>