



Integrated Management of Foliar Diseases of Mungbean (*Vigna radiata* L.) Under Natural Field Conditions

Ashok Kumar Meena ^{a*}, Neelam Geat ^a
and Vinay Kumar Kardam ^a

^a Agricultural Research Station, Mandore (Agriculture University), Jodhpur, 342304, Rajasthan, India.

Authors' contributions

This work was carried out in collaboration among all authors. Author AKM designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors NG and VKK managed the analyses of the study as well as managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jsrr/2024/v30i102434>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/122330>

Original Research Article

Received: 10/07/2024
Accepted: 12/09/2024
Published: 19/09/2024

ABSTRACT

Aims: To determine the efficacy of chemicals/bioagents against foliar diseases of mungbean.
Study Design: Randomized Block Design (Field experiments)
Place and Duration of the Study: Field experiments were conducted at Agricultural Research Station, Mandor, Jodhpur during the *Kharif* 2022 & *Kharif* 2023
Methodology: Different novel fungicides/bioagents were applied against foliar diseases of mungbean. Ten treatments *viz*; soil application with *Trichoderma harzianum* @ 5kg/ha enriched in 100 kg of FYM., seed treatment with *Trichoderma harzianum* @ 10g/kg seed, foliar spray of

*Corresponding author: E-mail: akmeenapathology@gmail.com;

Cite as: Meena, Ashok Kumar, Neelam Geat, and Vinay Kumar Kardam. 2024. "Integrated Management of Foliar Diseases of Mungbean (*Vigna Radiata* L.) Under Natural Field Conditions". *Journal of Scientific Research and Reports* 30 (10):88-94. <https://doi.org/10.9734/jsrr/2024/v30i102434>.

Trichoderma harzianum @ 2g/l, foliar spray of carbendazim 50% WP @ 0.1%, foliar spray of difenoconazole 25 EC @ 0.1 %, foliar spray of tebuconazole 250 EC @ 0.1%, foliar spray of tebuconazole 50% + trifloxystrobin 25% WG @ 0.1%, foliar spray of pyraclostrobin 133g/l+ epoxiconazole 50 g/l w/w SE @ 0.15%, foliar spray of azoxystrobin 18.2 % + difenoconazole 11.4 % SC @ 0.1%, and control (untreated) were taken in study. The diseases assessment was done after 10 days of spraying and percent disease incidence was calculated.

Results: Foliar spray of tebuconazole 50% + trifloxystrobin 25% WG @ 0.1% was most effective in reducing the anthracnose PDI (19.41%) and cercospora leaf spot (18.26%) followed by foliar spray of azoxystrobin 18.2 % + difenoconazole 11.4 % SC @ 0.1% (24.36 and 21.05 %).

Conclusion: Among all the tested treatments, foliar spray of tebuconazole 50% + trifloxystrobin 25% WG @ 0.1% was most effective followed by foliar spray of azoxystrobin 18.2 % + difenoconazole 11.4 % SC @ 0.1% in managing both the diseases.

Keywords: Mungbean; *Colletotrichum* sp.; *Cercospora* sp.; fungicides; bio agents.

1. INTRODUCTION

Mungbean or green gram [*Vigna radiata* (L.) R. Wilczek] is a nutrition-rich, short-duration, belonging to the family *Leguminosae* and warm season grain legume which is cultivated in several countries across Asia, East Africa and Australia. In India, it ranks third in importance amongst pulse crops, after pigeon pea and chickpea [1] is believed to have originated in India or the Indo-Burmese region. Mungbean is extensively cultivated in southern and eastern Asia and has a crucial dietary source of protein and essential nutrients. It provides 26 per cent protein, 51 per cent carbohydrates, 10 per cent moisture, 4 per cent minerals, and 3 per cent vitamins. Lysine is an important amino acid that is often lacking in cereal proteins, yet mungbean seeds are rich in it and used as fodder as well as dried seeds, this crop is a valuable source of food for humans. It is often grown in rotation with cereals to enhance soil fertility through its nitrogen-fixing ability. 90 per cent of the world's mungbean crop is produced in Asia, with India representing the leading producer, accounting for almost 50 per cent of the crop's global output while also consuming nearly all of it. The area, production and productivity of mungbean in Rajasthan are 2560 ha, 905 tonnes and 354 kg/ha, respectively [2]. However, mungbean cultivation faces numerous challenges, including diseases caused by fungi, bacteria, viruses, nematodes, and abiotic stresses.

“Among the fungal diseases, anthracnose caused by *Colletotrichum gloeosporioides* (Penz.) Penz. & Sacc. is one of the economically important disease. The anthracnose disease of mungbean was first reported in India from Jorhat of Assam state in 1951. The disease has been reported from all major mungbean growing states

of India in mild to severe form and in tropical and subtropical areas causes considerable damage by reducing seed quality and yield” [3]. “It is a seed-borne disease can lead to complete yield loss under favourable conditions when infecting susceptible cultivars. The average seed yield loss of 40.18 per cent and stalk yield loss of 46.90 per cent was noticed due to anthracnose of mungbean” [4]. “Another most important disease is, cercospora leaf spot (CLS) caused by the fungus *Cercospora canescens* Illis & Martin is a serious disease in mungbean (*Vigna radiata* (L.) Wilczek), The disease was reported first time in Delhi, India and disease can reduce seed yield by up to 50% and can be occurred in all parts of the humid tropical areas of Asia and many other countries and is prevalent in all parts of humid tropical areas of India, Bangladesh, Indonesia, Malaysia, Philippines, and Thailand. The disease causes qualitative and quantitative losses up to 96 per cent under natural epiphytotic conditions” [5-7]. Therefore, the current study aims to investigate the use of biocontrol agents in combination with fungicides under field conditions to develop an effective management strategy.

Thus, the primary objective of this study is to evaluate the efficacy of bioagents, both alone and in combination with fungicides, in various application modes. This approach aims to develop an integrated and effective management strategy to mitigate foliar disease impacts and reduce grain yield losses in mungbean.

2. MATERIALS AND METHODS

2.1 Details of the Experiment

The field experiment was conducted at the Agricultural Research Station, Mandore,

Table 1. Rating Scale for mungbean anthracnose disease

Scale	Description
0	No symptoms on leaves.
1	Small size lesions covering 1% or less of leaf area
3	Small size lesions covering 1-10% of leaf area.
5	Lesions size big but not coalescing covering,11-25% of the leaf area.
7	Lesions on leaves covering 26-50% of leaf area. Cankers on stem and pod infection.
9	Lesions on leaves covering 51% or more of leaf area. Defoliation of leaves, deep cankers on stem and pods, blighting of plant occurs.

Table 2. Rating Scale for mungbean cercospora leaf spot disease

Scale	Description
1	Small pinhead size brown lesions on leaf covering less than 1% leaf area
3	Small round, brown lesions on leaf covering 1-10% of leaf area
5	Lesions on leaves bigger, covering 1-25% of leaf area
7	Lesions enlarging and coalescing covering 26-50% of leaf area, shot-hole symptoms produced
9	Lesions coalescing 51%or more of leaf area, shot hole symptoms produced, lesions also on pods

Agriculture University, Jodhpur during the *Kharif* season 2022 and 2023, using mungbean variety GM-4. All the bio agents and fungicides were obtained from Agricultural Research Station, Mandor and complete package and practices were followed during entire crop season. Bioagent treatments were applied at the time of sowing, and fungicidal treatments were administered at the first appearance of the disease. A control plot was maintained for each replication without any fungicidal treatments. Observations on the intensity of anthracnose and cercospora diseases were recorded 15 days after the last spraying. Ten plants were randomly selected and labelled from each plot to assess disease intensity. The disease intensity was recorded by observing three trifoliate leaves from the basal, middle, and upper portions of the plants. These selected plants were graded using a 0-9 disease rating scale for anthracnose (Table 1) and 1-9 disease rating scale for cercospora (Table 2), based on the percentage area of leaves infected by the pathogen [8]. The seed yield (kg/ha) was also recorded.

The percent disease intensity (PDI) for both the disease was calculated by using the formula given by Mayee and Datar [9].

$$\text{Percent Disease Intensity (PDI)} = \frac{\text{Sum of numerical rating}}{\text{No.of leaves observed} \times \text{Maximum rating}} \times 100$$

3. RESULTS AND DISCUSSION

The results of this study demonstrate the significant impact of integrated management of anthracnose (*Colletotrichum lindemuthianum*) and cercospora leaf spot (*Cercospora canescens*) in mungbean crops. The field experiment's outcomes, as presented in Table 3, clearly indicated that the disease severity was significantly low in all the treated plots compared to the unsprayed control plot. The treatment with tebuconazole 50% + trifloxystrobin 25% WG at 0.1% was the most effective, resulting in the lowest disease intensities for both anthracnose (19.41%) and cercospora leaf spot (18.26%). This efficacy can be attributed to the synergistic action of tebuconazole, a triazole fungicide known for its broad-spectrum activity, and trifloxystrobin, a strobilurin fungicide that inhibits spore germination and mycelial growth. The combination of azoxystrobin 18.2% + difenoconazole 11.4% SC at 0.1% also showed substantial effectiveness, with disease intensities of 24.36% for anthracnose and 21.05% for cercospora leaf spot. azoxystrobin, another strobilurin fungicide, provides broad-spectrum control, while difenoconazole, a triazole, further enhances the treatment's efficacy. The untreated control plots showed the highest disease intensities, with 39.59% for anthracnose and 36.62% for cercospora leaf spot, and the lowest seed yield (1384 kg/ha).

Table 3. Integrated Management of Foliar Diseases of Mungbean

Treatment	Percent Disease Intensity (%)						Yield (Kg/ha)		
	Anthracnose			Cercospora leaf spot			2022	2023	Pooled
	2022	2023	Pooled	2022	2023	Pooled			
T ₁ Soil application with <i>Trichoderma harzianum</i> @ 5kg/ha enriched in 100 kg of FYM	31.33 (34.01) *	36.00 (36.85)	35.43	24.00 (29.24)	28.00 (31.91)	30.57	1023	975	999
T ₂ Seed treatment with <i>Trichoderma harzianum</i> @ 10g/kg seed	28.67 (32.33)	32.00 (34.42)	33.38	20.67 (26.75)	24.00 (29.28)	28.02	1043	1010	1026
T ₃ Foliar Spray of <i>Trichoderma harzianum</i> @ 2g/l	32.67 (34.70)	37.33 (37.66)	36.18	26.00 (30.61)	30.67 (33.62)	32.11	1000	781	890
T ₄ Foliar spray of carbendazim 50% WP @ 0.1%	22.00 (27.96)	28.00 (31.91)	29.93	17.33 (24.22)	21.33 (27.49)	25.86	1123	1063	1093
T ₅ Foliar spray of difenoconazole 25 EC @ 0.1 %	24.67 (29.58)	30.67 (33.62)	31.60	18.67 (25.34)	22.67 (28.36)	26.85	1095	1038	1067
T ₆ Foliar spray of tebuconazole 250 EC @ 0.1%	20.67 (26.93)	25.33 (30.21)	28.57	15.33 (22.94)	17.33 (24.57)	23.75	1153	1148	1151
T ₇ Foliar spray of tebuconazole 50% + trifloxystrobin 25% WG @ 0.1%	8.00 (16.35)	14.67 (22.47)	19.41	9.33 (17.53)	10.67 (18.99)	18.26	1413	1355	1384
T ₈ Foliar spray of pyraclostrobin 133g/l+ epoxiconazole 50 g/l w/w SE @ 0.15%	16.00 (23.47)	24.00 (29.33)	26.40	10.67 (18.95)	13.33 (21.37)	20.16	1213	1261	1237
T ₉ Foliar spray of azoxystrobin 18.2 % + difenoconazole 11.4 % SC @ 0.1%	14.67 (22.22)	20.00 (26.49)	24.36	14.67 (22.45)	11.33 (19.66)	21.05	1293	1289	1291
T ₁₀ Untreated control	36.00 (36.87)	45.33 (42.32)	39.59	32.00 (34.41)	39.33 (38.83)	36.62	971	571	771
SE(m) ±	2.09	1.14	2.39	2.28	1.20	2.58	55.34	61.12	58.30
CD at 5%	6.23	3.41	6.86	6.80	3.56	7.41	164.44	181.61	236.51
CV (%)	12.78	6.10	9.61	15.70	7.59	12.03	8.46	10.09	9.25

*Figures in parenthesis indicate angular transformation values

Table 4. Effect of Integrated Management of Foliar Diseases on economics of Mungbean

Treatment	Gross Return (₹/ha)			Net Return (₹/ha)			B:C Ratio			
	2022	2023	Mean	2022	2023	Mean	2022	2023	Mean	
T ₁	Soil application with <i>Trichoderma harzianum</i> @ 5kg/ha enriched in 100 kg of FYM	79334	83441	81387	52834	56941	54887	2.99	3.15	3.07
T ₂	Seed treatment with <i>Trichoderma harzianum</i> @ 10g/kg seed	80885	86436	83660	55435	60986	58210	3.18	3.40	3.29
T ₃	Foliar Spray of <i>Trichoderma harzianum</i> @ 2g/l	77550	66838	72194	51800	41088	46444	3.01	2.60	2.80
T ₄	Foliar spray of carbendazim 50% WP @ 0.1%	87089	90972	89030	60789	64672	62730	3.31	3.46	3.39
T ₅	Foliar spray of difenoconazole 25 EC @ 0.1 %	84917	88832	86875	58717	62632	60675	3.24	3.39	3.32
T ₆	Foliar spray of tebuconazole 250 EC @ 0.1%	89415	98246	93830	63215	72046	67630	3.41	3.75	3.58
T ₇	Foliar spray of tebuconazole 50% + trifloxystrobin 25% WG @ 0.1%	109578	115961	112770	79978	86561	83270	3.70	3.94	3.82
T ₈	Foliar spray of pyraclostrobin 133g/l+ epoxiconazole 50 g/l w/w SE @ 0.15%	94068	107916	100992	66618	80366	73492	3.43	3.92	3.67
T ₉	Foliar spray of azoxystrobin 18.2 % + difenoconazole 11.4 % SC @ 0.1%	100272	110313	105292	71672	81713	76692	3.51	3.86	3.68
T ₁₀	Untreated control	75301	48866	62084	50001	23566	36784	2.98	1.93	2.45

The economic analysis (Table 4) revealed that the foliar spray of tebuconazole 50% + trifloxystrobin 25% WG at 0.1% not only controlled the diseases effectively but also provided the highest benefit-cost (BC) ratio of 3.82. This indicates that the investment in this fungicidal treatment resulted in substantial economic returns due to the improved seed yield and reduced disease losses. The azoxystrobin 18.2% + difenoconazole 11.4% SC at 0.1% treatment also proved to be cost-effective, with a BC ratio of 3.68, demonstrating a favourable balance between the cost of treatment and the economic benefits of higher yields and lower disease intensity. The untreated control had the lowest (2.45) BC ratio. The observed economic benefits further justify the adoption of these fungicidal treatments by farmers, promoting sustainable and profitable mungbean cultivation.

Similarly, [10] reported that Carbendazim completely inhibited the mycelial growth of *C. canescens* at 200 ppm concentration under *in vitro* and bio-agent, *Trichoderma viride* was also found effective against the pathogen [11] also mentioned that least percent disease intensity of anthracnose was reported with fungicidal seed treatment with carbendazim + spraying with propiconazole followed by spraying of propiconazole alone. Combine seed treatment of carbendazim + *T. viride* also exhibited encouraging result in reducing disease intensity and higher seed yield and BC ratio. Similar outcomes have been accounted by Rajashree et al [12] that “among the combi fungicides evaluated, cent per cent inhibition was recorded in tricyclozole 18% + mancozeb 62%WP and carbendazim 12% + mancozeb 63% WP at 0.3 per cent concentration. In case of bioagents screened against *C. truncatum*, the highest mycelial inhibition was found in the *T. viride* strain (Tv- 29) followed by strain Tv- 1 and Tv- 10. While among the bacterial bioagents, *B. subtilis* strain (Bs-21) and *P. fluorescens* strain (Pf-26) showed highest mycelial inhibition”.

An observation was made [13] that “a combination of seed treatment with thiamethoxam at 4 g kg⁻¹, carboxin at 2 g kg⁻¹ and Pusa 5SD (*Trichoderma virens*) at 4 g kg⁻¹ followed by simultaneous foliar sprays of thiamethoxam 0.02% and carbendazim 0.05% at 21 and 35 days after sowing resulted in the highest seed germination and grain yield in mungbean with the lowest intensities of cercospora leaf spots and other diseases. Several researchers have been reported

effectiveness of chemicals in the past investigations” [14-17].

4. CONCLUSION

In conclusion, the field experiment demonstrated that the management approach using specific fungicide treatments significantly reduced the severity of anthracnose and cercospora leaf spot diseases in mungbean. The most effective and economically beneficial treatment was the foliar spray of tebuconazole 50% + trifloxystrobin 25% WG at 0.1%, followed closely by the azoxystrobin 18.2% + difenoconazole 11.4% SC at 0.1%. These treatments not only minimized disease intensity but also maximized seed yield and economic returns. The study demonstrates that effective fungicidal treatments not only reduce disease severity in mungbean but also enhance economic returns, making them a crucial component of integrated disease management strategies. The results provide a strong basis for recommending these treatments to mungbean farmers for sustainable crop production.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

ACKNOWLEDGEMENTS

Authors are thankful to Zonal Director Research of Agricultural Research Station, Mandore, Agriculture University, Jodhpur which was provided the facilities for conducting experimental trails.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Pratap A, Gupta S, Rathore MTB, Singh CM, Prajapati U, Singh P, Singh Y, Kumari G. The Beans and the Peas From Orphan to Mainstream Crops. 2021;1-32
2. Anonymous. Available: <https://rajas.rajasthan.gov.in/Ind ex.aspx>, Directorate of Economics & Statistics, Rajasthan 2021-22. 2022; Accessed 11th September 2024.

3. Chaudhari KA, Gohel NM. Management of anthracnose disease of mungbean through new fungicidal formulations. *Journal of Pure and Applied Microbiology*. 2016;10(1):691-696
4. Kulkarni SA. Epidemiology and integrated management of anthracnose of green gram. M.Sc. (Agri.) Thesis submitted to UAS, Dharwad, Karnataka; 2009; Accessed 11th September 2024.
5. Kumar N, Kumar S, Prajapati S, Maurya S. Cercospora leaf spot disease of green gram and its management: A review. *Journal of Pharmacognosy and Phytochemistry*. 2020;9(1):1574-1576
6. Chankaew S, Somta P, Sorajjapinun W, Sirinivas P. Quantitative trait loci mapping of Cercospora leaf spot resistance in mungbean, *Vigna radiata* (L.) Wilczek. *Mol Breeding*. 2011;(28):255–264. Available: <https://doi.org/10.1007/s11032-010-9478-1>
7. Kaur A, Gupta VP, Singh RB. Chemotherapy- an efficient tool to control Cercospora leaf spot in mungbean. *J Mycol. Pl. Pathol*. 2004;34:515-516.
8. Mayee CD, Datar VV. "Phytopathometry" Technical Bulletin-I, Marathwada Agricultural University, Parbhani, India. 1986;146.
9. Prasad D, Gupta K, Singh VP. Management of Cercospora Leaf Spot of Mungbean [*Vigna radiata* (L.) Wilczek] using Fungicides and Host Resistance in Bundelkhand Region of Uttar Pradesh, India. *Legume Research: An International Journal*. 2024;47(2):318.
10. Raghuvanshi RS, Chandra S, Singh A, Singh V, Pandey S, Raghuvanshi VV. Management of Cercospora Leaf Spot in Mungbean by use of Botanicals, Fungicides and Bio-agents. *Biological Forum – An International Journal*. 2022;14 (4):276-287.
11. Misal DM, Khaire PB, Misal MR, Hingole DG. Integrated evaluation of fungicides, botanicals and bioagents against anthracnose of mungbean on natural field condition. *International Journal of Chemical Studies*. 2019;7(2):1975-1978.
12. Rajashree G, Patil MB, Aswathanaryana DS, Mallikarjun K, Sreenivas AG. *In vitro* evaluation of different fungicides against *Colletotrichum truncatum* causing anthracnose of green gram. *International Journal of Chemical Studies*. 2019;7(5): 3295-3299.
13. Dubey SC, Singh B. Integrated management of major diseases of mungbean by seed treatment and foliar application of insecticide, fungicides and bioagent. *Crop Protection*. 2013;47:55-60.
14. Patel MK, Ranganathswamy Math. Evaluation of mix fungicides against *Colletotrichum truncatum* causing anthracnose of black gram. *The Pharma Innovation Journal*. 2023;12(9):2220-2223.
15. Purushotham P, Rakholiya KB, Vanani KD. Effectiveness of fungicides against *Colletotrichum lindemuthianum* causing anthracnose of green gram [*Vigna radiata* (L.) Wilczek]. *Legume Research*; 2023. Available: <https://arccjournals.com/journal/legume-research-an-international-journal/LR-5137>
16. Chaudhari KA, Gohel NM. Management of anthracnose disease of mungbean through new fungicidal formulations. *J. Pure Appl. Microbiol*. 2016;10(1):691-696.
17. Singh S, Prasad D, Singh VP. Evaluation of fungicides and genotypes against anthracnose disease of mungbean caused by *Colletotrichum lindemuthianum*. *International Journal of Bio-resource and Stress Management*. 2022;13(5):448-453.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/122330>