



# Investigation of the Ergonomics and Workload of Farm Women during in Planting and Weeding Operations

Naveendra Kumar Patel <sup>a++\*</sup>, Shubham Singh <sup>b</sup>  
and Utkarsh Dwivedi <sup>c#</sup>

<sup>a</sup> K.V.K. Sheikhpura, Bihar Agricultural University, Sabour, Bhagalpur, Bihar, India.

<sup>b</sup> Department of Farm Machinery and Power Engineering, VIAET, SHUATS, Prayagraj, Uttar Pradesh, India.

<sup>c</sup> College of Agricultural Engineering and Post-Harvest Technology, Central Agricultural University, Ranipool, Gangtok, Sikkim, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jsrr/2024/v30i62079>

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/117003>

**Original Research Article**

**Received: 14/03/2024**

**Accepted: 17/05/2024**

**Published: 21/05/2024**

## ABSTRACT

The study was done to determine the anthropometric, physical, physiological, and workload characteristics of women operators (N=7) during manual sowing and weeding activities for several medicinal plants. The planting and weeding activities were done using a shovel and a hand hoe. Anthropometric measurements were taken for women aged 20 to 50. The physiological fluctuations in chosen participants' energy and strength during different agricultural operations, as well as the

<sup>++</sup> Technical Assistant;

<sup>#</sup> Research Scholar;

<sup>\*</sup>Corresponding author: E-mail: [pnaveendra7378@gmail.com](mailto:pnaveendra7378@gmail.com);

**Cite as:** Patel, N. K., Singh, S., & Dwivedi, U. (2024). Investigation of the Ergonomics and Workload of Farm Women during in Planting and Weeding Operations. *Journal of Scientific Research and Reports*, 30(6), 615–622. <https://doi.org/10.9734/jsrr/2024/v30i62079>

implications in body part discomfort score, were evaluated. The average resting and working heart rates, EER, OCR, and BPDS are all measured to assess physiological characteristics. Female responders reported an average resting heart rate of 78.91 bpm for weeding and 81.46 bpm for planting. Planting and weeding had an average working heart rate of 100.85 and 97.14 (bpm, respectively). The average EER and OCR values for the relevant procedures were 6.35 and 5.31 (kJ/min) and 0.52 and 0.45 (l/min), respectively. The BPDS levels of the chosen participants were 19.5 and 21.4, respectively. The data imply that the workers were more compatible and comfortable when weeding using a hand hoe.

*Keywords: Energy expenditure; oxygen consumption rate; pulse rate and workload; weeding; energy expenditure.*

## 1. INTRODUCTION

Indian farmers are increasingly concerned with properly carrying out different farm activities to increase production [1,2]. They prioritize agricultural output over safety and comfort [3]. The man-machine interaction should be harmonic enough to provide safe and dependable horticultural operations while increasing farmers' working efficiency [4]. Since ergonomics are employed in agriculture to analyze the performance of the working operators undertaking any agricultural process, it may be readily useful in horticulture operations as well [5].

An ergonomic assessment is a technique for determining the energy expenditure of labor, its physiological cost, the method's appropriateness for farmworkers, and how long they can work without becoming weary [6]. Every farm job includes some drudgery, which causes physical and mental strain [7]. Women generally face greater drudgery than men [8]. The necessity for ergonomics is critical to determining the daily working performance of respondents involved in drudgery-inducing agricultural tasks (Awasthi et al. 2020).

The final step in the ergonomic evaluation to examine the variations arising throughout the weeding operation is the assessment of the operators' anthropometrical, physiological, and body postural discomfort [9-12]. The current anthropometric data of agricultural workers may be relevant in the design and development of manually operated weeders [13-16]. The most common horticultural chores for most farms are planting and weeding. Typically, all farm activities begin with the actions listed above. As a result, ergonomic characteristics should be given appropriate weightage to guarantee the safety and comfort of working operators [17,18,19].

As a result, the current findings stress the ergonomics considerations of female farm operators in horticulture to provide a safe and adaptive environment for working operators engaged in planting and weeding activities involved in medicinal plant production.

## 2. MATERIALS AND METHODS

### 2.1 Method for Implementing the Experimental Strategy in the Ergonomic Study

The study was conducted at Krishi Vigyan Kendra in Sheikhpura, Bihar, India. The selected respondents' anthropometric characteristics were assessed at the start of the investigation. Later, the subject's were permitted to seed medicinal plants such as tulsi and lemongrass using a shovel and hand hoe at the experimental location designated in the field, with a row-to-row spacing of 1.3 m and a period of 30 minutes. Fifteen female respondents from each age group (20-50 years old) were chosen at random based on their previous health history.

At the start of the experimental trials, all anthropometric measures were obtained with caution using an anthropometer, measuring tape, and steel scale, and different bodily characteristics of workers of all ages were recorded. They were then allowed to carry out the excavating and planting operations using the equipment [20-22].

The physiological observations of the individual participants were obtained before and during the procedure. Measurements of their resting and working pulse rates were taken and after five minutes of work started. The additional variables, such as OCR, EER, and BPDS, were then estimated based on the parameters given above.

A similar approach was used for weeding around the therapeutic plants.

## 2.2 Physiological and Psychological Characteristics of the Selected for farm Women Participants

### 2.2.1 Body mass index

The BMI is defined as the body mass divided by the square of the body height. The relation of BMI is given as.

$$BMI \left( \frac{kg}{m^2} \right) = \frac{Weight}{(Height)^2}$$

### 2.2.2 Lean body mass

Lean body mass is a component of body composition, calculated by subtracting body fat weight from total body weight. Hume's Equations of LBM for female respondents is given as follows.

$$\text{For female: } LBM = (0.29569 \times W) + (0.41813 \times H) - 43.2933$$

### 2.2.3 Basal metabolic rate

Basal metabolic rate is the number of calories required to keep your body functioning at rest, also known as the metabolism. It is related to body mass, age, weight and height. It is also affected by gender. According to Harris Benedict's Equations, BMR for female respondents is represented as follows.

$$\text{For female: } BMR = 655.1 + (9.563 \times W) + (1.850 \times H) - (4.676 \times \text{age in years})$$

Where,

$$W = \text{Body weight in kg}$$

$$H = \text{Body height in cm}$$

### 2.2.4 Pulse rate

The pulse rate was monitored and recorded using the heart rate monitor and pulse oximeter.

It was recorded before and after the start of the planting and weeding. The corresponding strength data is an illustration of materials utilized during the study in the Table 1 given below. Specifications and working features of the mentioned weeding implements are presented in Table 2.

### 2.2.5 Energy Expenditure Rate (EER)

Varghese et al., [23] proposed the following method for determining and estimating EER in their research of workers.

$$EER = 0.159 \times \text{Average heart rate} - 8.72 \text{ (KJ/min)}$$

### 2.2.6 Oxygen Consumption Rate (OCR)

Oxygen Consumption Rate was computed using the heart rate data that the operator has previously recorded. The following is an equation that depicts OCR [24]:

$$OCR \text{ (L/min)} = 0.0114 \times HR - 0.68$$

### 2.2.7 Body Part Discomfort Score (BPDS)

Techniques from Corlett et al. [25] were used to measure the localized discomfort. This approach divides the individuals' bodies into 27 regions. To discourage a response marking only one body region, each was assigned a separate number. The selected individuals were asked to list all body locations that caused discomfort, beginning with the most severe and progressing in decreasing order until no more places were mentioned. The aforesaid conclusions were determined by the following connection, which is listed below.

$$BPDS = \sum X_i \times S \text{ (3.40)}$$

Where,

$$X_i = \text{Number of body parts}$$


$$S = \text{Discomfort score (is this on a scale of 6 to 1?)}$$

**Table 1. An illustration of materials (equipment) utilized during the study**

Characteristics	Variables	Equipment's / Tools
Anthropometric	To measure body dimensions	Anthropometer, steel scale, measuring tape
Physical variables	Weight	Weighing balance
Physiological responses	Pulse rate, blood pressure	Heart rate monitor, Pulse oximeter

**Table 2. Specifications and working features of the mentioned weeding implements**

Shovel		Hand hoe (khurpa)	
Overall length (mm)	735	Overall length (mm)	320
Width of cutting edge (mm)	315	Width of cutting edge (mm)	105
Weight (kg)	2.5	Weight (kg)	1.1
Working depth (mm)	155	Working depth (mm)	88



**Fig. 1. Region for evaluating body part discomfort score**

Anthropometric data of chosen participants were collected using a measuring anthropometer in total resting state. Seven subjects were selected from agricultural farms of different selected in age subject (Table 3 and Table 4).

In the current study, the meteorological conditions prevalent during planting and weeding operations, such as average temperature, humidity, and quantity of sunshine, were also examined throughout replications of the

**Table 3. Anthropometric information regarding the chosen participants**

S.No.	Anthropometric & strength data	Age of women operators (years)						
		20	25	30	35	40	45	50
1	Weight(kg)	47	45	49	52	56	61	59
2	Stature (cm)	156	151	145	158	161	166	163
3	Elbow height(cm)	98	93	89	96	101	105	96
4	Olecranon height(cm)	95	89	86	94	97	98	91
5	Illiocrystable height(cm)	88	81	80	85	87	89	84
6	Illiospinal height(cm)	143	146	136	141	148	153	145
7	Knee height(cm)	49	51	42	48	49	53	50
8	Arm reach(cm)	75	71	62	72	76	78	75
9	Vertical reach(cm)	192	185	181	190	194	198	186
10	Hand length(cm)	63	65	63	67	69	72	69
11	Head length(cm)	17	18	17	19	18	19	18
12	Foot length(cm)	22	21	20	22	23	24	22

(Measurement is taken in cm unless otherwise specified)

**Table 4. Physical traits possessed by operators**

S. No.	Physical characteristics	Age (years)							Average
		20	25	30	35	40	45	50	
1.	Age (years)	20	25	30	35	40	45	50	
2.	Height (cm)	156	151	145	158	161	166	163	157.14
3.	Weight (kg)	47	45	49	52	56	61	59	52.71
4.	BMI (kg/m <sup>2</sup> )	19.31	19.74	23.31	20.83	21.60	22.14	22.21	21.30
5.	LBM	35.83	33.15	31.82	38.15	40.58	44.15	42.31	38.00
6.	BMR	1299.641	1247.89	1251.66	1281.02	1301.44	1335.12	1287.07	1286.26

**Table 5. Enumer Operating conditions during different farm operations**

S. No.	Months	Planting	Weeding
		July (2022)	August (2022)
1.	Average temperature, °C	36	39
2.	Average humidity, %	71	75
3.	Average sunshine, hours	8.3	7.7

procedure taken in corresponding months, as illustrated in the Table 5 given.

### 3. RESULTS AND DISCUSSION

#### 3.1 The influence physiological characteristics of selected subjects performing variance farm operators

In all agricultural activities, the average resting heart rate values of the respondents, who were 20, 25, 30, 35, 40, 45, and 50 years old, were recorded as 72, 76, 73, 79, 76, 78, and 80, respectively.

Similarly, the individuals' average working heart rates throughout planting and weeding activities were 101.86 and 96.86 bpm, respectively. As a result, the subjects were more comfortable and familiar with weeding with a hand hoe since it is a light workload, as opposed to weeding with a shovel, which requires more effort and entails moderate work [26].

Similarly, the average EER for respondents using a shovel to dig and sow herbal plants (planting)

was 7.48 kJ/min, while the average EER for using a hand hoe to carry out weeding operations was 6.68 (kJ/min), as shown in Table 7.

The recent showed that more energy demand was doing digging with shovel for weeding. However energy expenditure during weeding with hand hoe was less due to its lightweight Similarly, the average OCR for the selected operators during planting and weeding operation were 0.48 and 0.42 (L/min) respectively as enumerated in the Table 8. Therefore, the subjects were more comfortable and quite familiar while weeding operation by hand hoe as it involves effort and ease.

This provides less oxygen consumption throughout the operation than a shovel, which requires maximal oxygen uptake since pain and tiredness occur while operating a shovel. The latter procedure requires continual arm action to elevate and lower the implement, hence the operators needed the most oxygen [26].

**Table 6. Assessment of the average pulse rate of the female operators in various farm operations**

S. No	Operations	Average pulse rate (bpm)							Average	
		Age (years)	20	25	30	35	40	45		50
1.	Planting operation		89	95	96	99	105	111	118	101.86*
2.	Weeding operation		84	90	88	95	101	109	111	96.86*

\*Significant at 0.05 level of significance

**Table 7. Assessment of average EER of the female operators in planting and weeding operations**

S. No	Operations	Average Energy Expenditure Rate (EER)(kJ/min) at varying age							
		Age (years)	20	25	30	35	40	45	50
1.	Planting operation	5.43	6.39	6.54	7.02	7.98	8.93	10.04	7.48*
2.	Weeding operation	4.64	5.59	5.27	6.39	7.34	8.61	8.93	6.68*

\*Significant at 0.05 level of significance

**Table 8. Assessment of average OCR of the female operators in sowing/planting and weeding operations**

S. No	Operations	Average Oxygen Consumption Rate (OCR, L min <sup>-1</sup> )							
		Age (years)	20	25	30	35	40	45	50
1.	Planting operation	0.33	0.4	0.41	0.45	0.52	0.59	0.67	0.48*
2.	Weeding operation	0.28	0.35	0.32	0.4	0.47	0.56	0.59	0.42*

\*Significant at 0.05 level of significance

**Table 9. Assessment of average BPDS of the female operators in various farm operations**

S.No	Operations	Average Body Part Discomfort Score (BPDS)							
		Age (years)	20	25	30	35	40	45	50
1.	Planting operation	19.04	20.4	22.44	22.44	22.44	23.12	24.48	22.05*
2.	Weeding operation	18.02	19.72	20.4	21.08	21.42	22.44	23.12	20.89*

\*Significant at 0.05 level of significance

### 3.2 Effect of Body Part Discomfort Score on the Performance of the Respondents

Table 9 shows the average BPDS values of the participants selected for planting and weeding activities, which were 22.05 and 20.89, respectively. Previous research found similar results [27].

### 3.3 Assessment of Physiological Responses of the Subjects

The age of farm women has a substantial impact on their heart rate, EER and OCR. At the 0.05 level of significance, physiological variables altered more substantially during planting than weeding. In terms of physiological reactions, female operators were more comfortable in a crouching posture, i.e., hand hoeing rather than digging cum replanting [28]. The pulse rate was found to be elevated among responders who used shovels due to the numerous efforts required and the implement's improper design. The operators' fitness difficulties have a direct impact on their heart rate when carrying out the digging and planting activity. As a result, the subjects were more comfortable and familiar with weeding with a hand hoe since it is a light workload, as opposed to weeding with a shovel,

which requires more effort and entails moderate work [26].

The current findings analyze whether respondents using a shovel consumed more oxygen than those using a hand hoe. Because the tiredness rate among shovel operators was rather high, oxygen consumption increased marginally as respondents' ages increased. As a result, the participants felt more at ease and comfortable when weeding with a hand hoe because it required little effort and was quite simple. This provides less oxygen consumption throughout the operation than a shovel, which requires maximal oxygen uptake due to pain and tiredness caused by using a shovel. The latter procedure requires continual arm movement to elevate and lower the implement, hence the operators needed the most oxygen [26].

The energy consumption rate was negligible when the operators used a hand hoe to weed rather than a shovel to seed. This might be due to the discomfort of continuing to work in a standing position while rising and elevating the arms, as well as the irregularity of using a shovel. Also, the current conditions during strong sunshine may be the cause of increased energy consumption. As a result, the participants used more energy during planting because controlling

a weighted shovel for digging demands significant effort, however for weeding, the operators were flexible and accustomed with using a hand hoe due to its lightweight [26].

### 3.4 Assessment of the Body Part Discomfort Score (BPDS) of the operators

The BPDS is significantly impacted by the age of farm women, with a greater variance in replanting operations compared to weeding activities at the 0.05 level of significance. The current investigation demonstrates that the bodily pain experienced was weariness in the knee and shoulder during the weeding operation, but it was an arm, knee, wrist, and shoulder during the sowing operation done by hand hoe due to the significant effort required by the arms to uproot the weeds. It became apparent that the hand hoe was well-designed, and its sharp blade required little effort to clear weeds. As a result, the research findings show that the hand hoe is more adaptive and stable when weeding in a sitting stance than managing a shovel in a standing position.

## 4. CONCLUSION

The respondents preferred to use a hand hoe over a shovel, which may be attributed to the former's consistency and ease of adaptation, as well as the latter's substantial weight, which made handling difficult. Furthermore, while weeding and planting, workers were compelled to frequently elevate and drop their arms in a bending position, which imposed additional strain on their shoulders. This generated physical pain and an increase in physiological variables.

## CONSENT

As per international standards or university standards, patient(s) written consent has been collected and preserved by the author(s).

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Pandey NG. Small Farmers in India: Challenges and Opportunities. Golden Research Thoughts. 2016;5.
2. Pandey S, Verma AK, Sirmour A, Gandhi I, Vishwavidyalaya K. Ergonomical studies on single row power weeder for rice crop. *croplandweed.com*. 2019;15:145–50.
3. Goel AK, Behera D, Behera BK, Mohanty SK, Nanda SK. Development and ergonomic evaluation of manually operated weeder for dry land crops. *Agricultural Engineering International: CIGR Journal*. 2008; Manuscript PM 08 009. Vol. X.
4. Sahoo L, Tanuja S, Argade S. Strengthening farm women perspective in quality seed production. Model Training Course "Strengthening Farm Women Perspective in Quality Seed Production. 2017;1–4.
5. Benos L, Tsaopoulos D, Bochtis D. A review on ergonomics in agriculture. Part I: Manual operations. *Applied Sciences (Switzerland)*; 2020.
6. Sam B. Ergonomic evaluation of paddy seeder and rotary weeder with women operators. *Proceedings 19th Triennial Congress of the IEA*. 2015;9–14.
7. Tiwari N, Upadhyay R, Dudi A. Adoption of Drudgery Reducing Tools among Farm Women. *Indian Research Journal of Extension Education*. 2021; 21:108–11.
8. Moharana G, Nayak J, Mhatre C, Rout PK. Assessment of Drudgery of Women in Agriculture. *Strengthening farm women perspective in quality seed production*. 2017;77.
9. Singh SK, Yadav RK, Kumar A. An Ergonomically Assessment of Manually Operated Maize Planter for Male Operators. *International Journal of Bio-resource and Stress Management*. 2023;14(6):872-7.
10. Fathallah FA. Musculoskeletal disorders in labor-intensive agriculture. *Applied Ergonomics*. 2010;41(6):738–43.
11. Gangopadhyay S, Dev S. Design and evaluation of ergonomic interventions for the prevention of musculoskeletal disorders in India. *Annals of Occupational and Environmental Medicine*. 2014;26.
12. Hemanth KG, Harshavardhan M, Ashok KA, Pavan KE, Ramana MV. Physiological evaluation of manual operated weeders and sprayers on farm use. *Agricultural Engineering International: CIGR Journal*. 2015;17:173–83.
13. Khogare DT, Sunita B. Anthropometric data of agricultural workers for suggesting dimensions of manually operated weeder. *Asian Journal of Home Science*. 2011;6:57–60.

14. Khogare DT, Borkar SB. Anthropometric data of agricultural workers for suggesting dimensions of manually operated weeder. 2011.
15. Mushobozi WL. Good agricultural practices (GAP) on horticultural production for extension staff in Tanzania: Training Manual. FAO GAP working paper series. 2010;13:101–10.
16. Murrell K. Ergonomics: Man in his working environment. Springer Science & Business Media. 2012.
17. Palega M, Rydz D. Work safety and ergonomics at the workplace an excavator. International Scientific Journal 'Trans Motoauto World'. 2018;29:25–9.
18. Karwowski W. Ergonomics: How to Design for Ease and Efficiency by Karl Kroemer, Henrike Kroemer, & Katrin Kroemer-Elbert 1994, 766 pages, \$86.00 Englewood Cliffs, NJ: Prentice-Hall ISBN 0-13-278359-2. Ergonomics in Design: The Quarterly of Human Factors Applications. 1997;5:33–5.
19. Kirkhorn MPH, Steven R, Earle-Richardson G, Banks CIE, Banks R. Ergonomic risks and musculoskeletal disorders in production agriculture: recommendations for effective research to practice. Taylor & Francis. 2010;15:281–99.
20. Saha PN. Practical use of some physiological research methods for assessment of work stress. Indian Association of Physiotherapists. 1976;4:9–13.
21. Awasthi VJ, Singh M, Mishra R, Chaudhary R, Pandey M, Parihar DS, Gautam S, Kumar D. Ergonomic and Workload Assessment in Weeding Operation Conducted for Wheat Crop for Male Respondents. Int J Curr Microbiol App Sci. 2019;8(09):2597-609.
22. Yadav R, Pund S. Development and ergonomic evaluation of manual weeder. Agricultural Engineering International: the CIGR Ejournal. 2007; Manuscript PM 07 022. Vol IX, October.
23. Varghese MA, Saha PN, Atreya N. A rapid appraisal of occupational workload from a modified scale of perceived exertion. Ergonomics. 1994;37:485–91.
24. Singh SP, Gite LP, Majumder J, Agarwal N. Aerobic capacity of farm women using sub-maximal exercise technique on tread mill. Agricultural Engineering International: the CIGR E Journal. 2008;10: Manuscript MES 08 001.
25. Corlett EN, Bishop RP. A technique for assessing postural discomfort. Ergonomics. 1976;19:175–82.
26. Swapnali B, Nilakshi B. Physiological Workload and Health Hazards of Tribal Women of Meghalaya Involved in Weeding Operation. International Journal of Current Microbiology and Applied Sciences. 2020;9(12):2589-95.
27. Kumar AA, Haribabu B, Rao AS, Someswararao C. Ergonomical evaluation of manually operated weeder under wet land condition. Scientific Research and Essays. 2013;8:249–55.
28. Panigrahi CMA. Managing stress at workplace. Journal of Management Research and Analysis. 2016;3(4):154-60.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/117003>